

Lund 1975

EVOLUTIONARY TRENDS IN THE ATRIPLEX
PROSTRATA GROUP OF SCANDINAVIA

BY MATS GUSTAFSSON

EVOLUTIONARY TRENDS IN THE ATRIPLEX PROSTRATA GROUP OF SCANDINAVIA

AV

MATS GUSTAFSSON

FIL. KAND., HLD



Akademisk avhandling som för avläggande av filosofie doktorsexamen vid Matematisk-Naturvetenskapliga Fakulteten vid Universitetet i Lund kommer att offentligen försvaras å föreläsningssalen vid Institutionen för Systematisk Botanik, torsdagen den 15 maj 1975 klockan 10.00.

Lund 1975

EXHIBITION OF THE AMERICAN ANTHROPOLOGICAL ARCHIVES

1894

1894

1894



Exhibition of the American Anthropological Archives
The American Anthropological Archives are the
collections of the American Anthropological Association
and the American Museum of Natural History. The
collections are housed in the American Anthropological
Archives, which are located in the American Museum of
Natural History, 200th Street, New York, N.Y. 10021.

The results are based on the following papers:

1. *Journal of the Royal Microscopical Society*, 1911, 31, 1-10.

2. *Journal of the Royal Microscopical Society*, 1912, 32, 1-10.

3. *Journal of the Royal Microscopical Society*, 1913, 33, 1-10.

4. *Journal of the Royal Microscopical Society*, 1914, 34, 1-10.

5. *Journal of the Royal Microscopical Society*, 1915, 35, 1-10.

6. *Journal of the Royal Microscopical Society*, 1916, 36, 1-10.

7. *Journal of the Royal Microscopical Society*, 1917, 37, 1-10.

8. *Journal of the Royal Microscopical Society*, 1918, 38, 1-10.

9. *Journal of the Royal Microscopical Society*, 1919, 39, 1-10.

10. *Journal of the Royal Microscopical Society*, 1920, 40, 1-10.

The following is a list of the names of the persons who have been associated with the work:

1. *Journal of the Royal Microscopical Society*, 1911, 31, 1-10.

2. *Journal of the Royal Microscopical Society*, 1912, 32, 1-10.

3. *Journal of the Royal Microscopical Society*, 1913, 33, 1-10.

4. *Journal of the Royal Microscopical Society*, 1914, 34, 1-10.

5. *Journal of the Royal Microscopical Society*, 1915, 35, 1-10.

6. *Journal of the Royal Microscopical Society*, 1916, 36, 1-10.

7. *Journal of the Royal Microscopical Society*, 1917, 37, 1-10.

8. *Journal of the Royal Microscopical Society*, 1918, 38, 1-10.

9. *Journal of the Royal Microscopical Society*, 1919, 39, 1-10.

10. *Journal of the Royal Microscopical Society*, 1920, 40, 1-10.

TO THE MEMORY OF GÖTE TURESSON
WHO INITIATED THE WORK

The following is a list of the names of the persons who have been associated with the work:

1. *Journal of the Royal Microscopical Society*, 1911, 31, 1-10.

2. *Journal of the Royal Microscopical Society*, 1912, 32, 1-10.

3. *Journal of the Royal Microscopical Society*, 1913, 33, 1-10.

4. *Journal of the Royal Microscopical Society*, 1914, 34, 1-10.

5. *Journal of the Royal Microscopical Society*, 1915, 35, 1-10.

6. *Journal of the Royal Microscopical Society*, 1916, 36, 1-10.

7. *Journal of the Royal Microscopical Society*, 1917, 37, 1-10.

8. *Journal of the Royal Microscopical Society*, 1918, 38, 1-10.

9. *Journal of the Royal Microscopical Society*, 1919, 39, 1-10.

10. *Journal of the Royal Microscopical Society*, 1920, 40, 1-10.



TO THE MEMORIAL OF JOHN THURGOOD

WHO TRAVELED THE WORLD

The thesis is based on the following papers:

- Gustafsson, M. 1973 a. Evolutionary trends in the *Atriplex triangularis* group of Scandinavia. I, Hybrid sterility and chromosomal differentiation. -- Bot. Notiser 126: 345--392.
- 1973 b. Evolutionary trends in the *Atriplex triangularis* group of Scandinavia. II. Spontaneous hybridization in relation to reproductive isolation. -- Bot. Notiser 126: 398--416.
- 1974. Evolutionary trends in the *Atriplex triangularis* group of Scandinavia. III. The effects of population size and introgression on chromosomal differentiation. -- Bot. Notiser 127: 125--148.
- 1975. Evolutionary trends in the *Atriplex prostrata* group of Scandinavia. IV. Taxonomy and morphological variation. -- Opera Bot. (in manuscript).

DELIMITATION OF THE *A. PROSTRATA* GROUP

The *A. prostrata* group comprises halophytic species with green, rhombic to hastate lower leaves and with bracteoles united up to the middle at most. In Scandinavia five taxa are recognized, viz. *A. glabriuscula* Edmondst., *A. longipes* Drej. ssp. *longipes*, *A. longipes* Drej. ssp. *praecox* (Hülph.) Turess., *A. prostrata* DC. ssp. *prostrata* and *A. prostrata* DC. ssp. *calotheca* (Rafn) M. Gust. comb. nov. All the taxa are annuals and diploids with the chromosome number $2n=18$.

MORPHOLOGY AND MORPHOLOGICAL VARIATION

The intrapopulation variation is particularly pronounced in *A. prostrata* ssp. *prostrata* and most populations are highly polymorphic in characters such as habit, height, size and shape of leaves and bracteoles. In the other taxa, a wide intrapopulation variation can be observed in some populations and a relatively narrow range of variation in others. In both *A. longipes* ssp. *longipes* and *A. prostrata* ssp. *prostrata* morphological differences between populations are rather small and the degree of over-

lapping extensive. In *A. glabriuscula*, *A. longipes* ssp. *praecox* and *A. prostrata* ssp. *calotheca* morphological differences exist between some populations but not between others. Many characters show a high degree of overlapping between taxa. The diagnostic characters are limited to the shape of leaves and bracteoles. However, other biological differences also exist, for instance rate of vegetative development, flowering time, length of the protogynous stage, floating capacity of the bracteoles enclosing the seed, ecological preferences, and population structure.

CULTIVATION EXPERIMENTS

Cultivation experiments have been carried out in all taxa, but primarily in *A. prostrata* ssp. *prostrata*. Characters used for distinguishing taxa within the polymorphic *A. prostrata* ssp. *prostrata*, such as height and leaf size but to some extent also habit and shape of leaves, are highly modificative.

HABITAT AND DISTRIBUTION

All the taxa inhabit seashores of different types, *A. prostrata* ssp. *prostrata* also occurring on waste ground. *A. glabriuscula* and *A. prostrata* ssp. *calotheca* are mainly distributed in the southwestern parts of Scandinavia, *A. longipes* ssp. *longipes* in the southern parts, *A. longipes* ssp. *praecox* in the Baltic area and in the northern parts of Norway, while *A. prostrata* ssp. *prostrata* is distributed throughout the whole of Scandinavia. The present patterns of distribution indicate that *A. longipes* ssp. *praecox* is of Arctic origin, while the other taxa have probably reached Scandinavia from the southwest.

POPULATION STRUCTURE

Most of the populations of *A. longipes* ssp. *longipes*, *A. prostrata* ssp. *prostrata* and ssp. *calotheca* investigated are made up of a

Digitized by the Internet Archive
in 2026

large number of individuals, about $2/3$ of these populations being large or very large, i.e. comprising more than 1,000 individuals. The majority of populations of *A. glabriuscula* are of medium size, some large, i.e. they comprise more than 300 individuals. By contrast, most populations of *A. longipes* ssp. *praecox* are small or very small the number of individuals usually varying between 1 and 300.

BREEDING SYSTEM

All the taxa are monoecious and highly self-fertile. *A. longipes* has a relatively long period of protogyny, while *A. glabriuscula* and *A. prostrata* are virtually not protogynous. Protogyny favours cross fertilization in *A. longipes*. In the other taxa the degree of self fertilization seems to be high, even dominant. The pollen grains are probably dispersed by the agency of wind.

GERMINATION

Experiments on germination indicate that seed samples of "pure" taxa germinate to a high degree (usually 60--100 %). However, for successful germination the seeds need at least a short period of dormancy, and in addition the temperature must fluctuate above and below zero.

FERTILITY AND MEIOSIS

The frequency of plants with reduced male fertility (less than 90 % stainable pollen) has been investigated in 6 populations of *A. glabriuscula*, 16 populations of *A. longipes* ssp. *longipes*, 14 populations of *A. longipes* ssp. *praecox*, 19 populations of *A. prostrata* ssp. *prostrata* and 6 populations of *A. prostrata* ssp. *calotheca*. The following results were obtained: 1. Plants with reduced fertility occur most commonly in populations of *A. longipes* ssp. *praecox* (about 20 % of the material), less commonly in the other taxa (6--9 %). 2. The frequency of plants with reduced fertility varies within populations of all taxa.

3. In *A. longipes* ssp. *praecox* plants with reduced fertility are commoner in small populations than in medium-sized to large ones.
4. In populations of *A. longipes* ssp. *longipes* and *A. prostrata* ssp. *prostrata* affected by introgression, the frequency of plants with reduced fertility is higher than in non-introgressive populations.

In all plants meiotically investigated, reduced fertility is to a great extent caused by the formation of bridges and laggards at anaphase I. Bridge-formation is probably due to heterozygosity for paracentric inversions.

REPRODUCTIVE ISOLATION AND HYBRIDIZATION

On the whole, the reproductive isolation between taxa is relatively small, but varies in different regions of Scandinavia. In the Baltic region, *A. longipes* ssp. *praecox* is ecologically isolated from *A. longipes* ssp. *longipes* and *A. prostrata* ssp. *calotheca*, and seasonal isolation exists between *A. longipes* and the other taxa. In W Scandinavia the seasonal isolation is only partial due to the fact that the flowering periods differ somewhat although overlapping occurs. In the northern parts of Norway the taxa flower simultaneously. Consequently hybridization and the occurrence of hybrid derivatives are relatively rare in the Baltic region, but relatively common in W Scandinavia and N Norway. Although reproductive isolation is rather weak the taxa remain relatively distinct.

The investigation of cultivated hybrid progenies derived from seeds produced by natural hybrids indicates the following: 1. Hybrid seeds usually germinate poorly compared with seed samples of "pure" taxa. 2. Plants with reduced fertility occur in all hybrid progenies, but this is most pronounced in *A. glabriuscula* x *A. prostrata* ssp. *prostrata*, and in *A. longipes* ssp. *longipes* x *A. prostrata* ssp. *prostrata*. 3. Morphological segregation towards the pa-

rent species is not necessarily combined with restoration of fertility.

INTROGRESSION

Introgression occurs between most taxa, but is most frequent between *A. longipes* and *A. prostrata* ssp. *prostrata*, and between *A. prostrata* ssp. *prostrata* and ssp. *calotheca*. This has lead to: 1. The transfer of morphological traits from one taxon to another. 2. An increased frequency of plants with reduced fertility. 3. In *A. prostrata* ssp. *prostrata* reduced fertility is commoner in F_1 progenies of crosses between non-introgressive populations than of crosses between introgressive populations.

CROSSING EXPERIMENTS

In the *A. prostrata* group crosses are easily performed, but seeds of intraspecific as well of interspecific crosses germinate poorly. The germination is low compared with that of seed samples of "pure" taxa. The majority of the hybrids are well-developed vegetatively and the seed-setting is good.

The aims of the crossing experiments have been to investigate: 1. Crosses between isolated populations within taxa, with special reference to morphological differences and geographical distance between the parent populations crossed in relation to hybrid fertility. 2. The mode and strength of internal barriers in crosses between taxa. 3. The relation between hybrid fertility and size of parent populations crossed. 4. The relation between chromosomal alterations and reduced hybrid fertility.

F_1 fertility in crosses within taxa

Interpopulational crosses have been carried out within *A. longipes* ssp. *longipes* (48 crosses), ssp. *praecox* (37) and *A. prostrata* ssp. *prostrata* (88). The following results were obtained: 1. Reduced fertility is common in crosses within *A. longipes* ssp. *praecox*, where about 80 % of the crosses have fertility values of less than 80 % stainable

pollen. Reduced fertility is less pronounced in crosses within *A. longipes* ssp. *longipes* and *A. prostrata* ssp. *prostrata*, where the corresponding figures are 15 % and 28 % respectively. 2. In *A. longipes* ssp. *longipes* and ssp. *praecox* little or no correlation exists between geographical distance separating parent populations and the fertility of their hybrid offspring. 3. In all three taxa there is virtually no correlation between hybrid fertility and morphological diversities between the parent populations crossed.

F_1 fertility in crosses between taxa

In some combinations the number of crosses is small, due to experimental difficulties. The results indicate, however, that the fertility of most crosses entailing *A. longipes* ssp. *praecox* is reduced (mean values about 70 % stainable pollen), while crosses between the other taxa are generally highly fertile (about 85—90 %). Baltic populations of *A. longipes* ssp. *praecox* have been crossed with populations of *A. longipes* ssp. *longipes* from the Baltic region, W Sweden and Denmark respectively. Reduced hybrid fertility is most pronounced in the Baltic region, where the taxa occur sympatrically. Selection for a crossing barrier has probably taken place in this region.

Fertility and morphological variation in F_2 progenies

Eighty-nine F_2 progenies were raised by selfing F_1 hybrids representing different crossing combinations at different levels of fertility. In most crosses between taxa there is a range from almost ^mcompletely fertile F_2 hybrids to hybrids with varying degrees of reduced fertility. Restoration of fertility seems to occur more readily in intraspecific than in interspecific crosses. Analysis of F_2 progenies indicates that most morphological characters are inherited independent of one another and that at least some traits are conditioned by more than one pair of genes.

The effects of population size on hybrid fertility

The degree of reduced fertility in crosses between populations of differ-

ent size has been investigated in intraspecific as well as in interspecific combinations. The results show that crosses with small populations of *A. longipes* ssp. *praecox* have a much higher frequency of reduced fertility than crosses between medium to large populations of ssp. *praecox* and of the other taxa.

Meiotic behaviour in hybrid plants

Meiosis has been investigated in representatives of 42 intraspecific as well as interspecific crosses. At metaphase I nine bivalents are regularly formed. No multivalents have been observed, but univalents were observed in two crosses. At anaphase I the number of cells with visible disturbances is often rather high, but these are exclusively of two types, viz. bridges and laggards. Usually single bridges have been observed, but two bridges have been observed occasionally. The correlation between the frequency of bridges formed at anaphase I and hybrid fertility is high ($r=0.72$), the lower the fertility, the higher the frequency of bridges. In the *A. prostrata* group all information indicates that formation of bridges at anaphase I is probably due to heterozygosity for paracentric inversions. Analysis of F_2 progenies indicates that in several crosses reduced fertility is caused by two small inversions rather than by a single large one.

INVERSION POLYMORPHISM

Investigations of fertility in natural populations, hybrid fertility and meiotic behaviour indicate that polymorphism for paracentric inversions is most pronounced in small populations of *A. longipes* ssp. *praecox*, less so in medium to large *praecox* populations and in *A. prostrata* ssp. *prostrata*, and is of little significance in the other taxa.

EVOLUTION

The extent and rate of intra- and interspecific differences in the *A. prostrata* group in Scandinavia is apparently to a great extent based on

evolution that took place before the last glaciation or in ice-free areas during the last glaciation. The present pattern of variation thus depends on the origin of the Scandinavian material and the different migratory routes, as well as on evolution, adaptation and gene flow within Scandinavia after the last glaciation, i.e. within the last 10,000 years.

